

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 September 2002 (19.09.2002)

PCT

(10) International Publication Number  
**WO 02/072774 A2**

(51) International Patent Classification<sup>7</sup>: C12N

(21) International Application Number: PCT/US02/07438

(22) International Filing Date: 6 March 2002 (06.03.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/275,009 12 March 2001 (12.03.2001) US  
60/284,152 17 April 2001 (17.04.2001) US

(71) Applicant: **LEXICON GENETICS INCORPORATED**  
[US/US]; 4000 Research Forest Drive, The Woodlands, TX  
77381 (US).

(72) Inventors: **WILGANOWSKI, Nathaniel, L.**; 505 Cypress Station #1906, Houston, TX 77090 (US). **NEPOM-NICHY, Boris**; 905 Cypress Station Dr. #1515, Houston, TX 77090 (US). **BURNETT, Michael, B.**; 1414 Corral Drive, Houston, TX 77090 (US). **HU, Yi**; 333 Holly Creek Ct. #203, The Woodlands, TX 77381 (US).

(74) Agents: **ISHIMOTO, Lance, K.** et al.; Lexicon Genetics Incorporated, 4000 Research Forest Drive, The Woodlands, TX 77381 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— without international search report and to be republished upon receipt of that report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



**WO 02/072774 A2**

(54) Title: NOVEL HUMAN TRANSPORTER PROTEINS AND POLYNUCLEOTIDES ENCODING THE SAME

(57) Abstract: Novel human polynucleotide and polypeptide sequences are disclosed that can be used in therapeutic, diagnostic, and pharmacogenomic applications.

WO 02/072774

PCT/US02/07438

NOVEL HUMAN TRANSPORTER PROTEINS AND  
POLYNUCLEOTIDES ENCODING THE SAME

The present application claims the benefit of U.S.  
5 Provisional Application Numbers 60/275,009 and 60/284,152,  
which were filed on March 12, 2001 and April 17, 2001,  
respectively. These provisional applications are herein  
incorporated by reference in their entirety.

## 1. INTRODUCTION

10 The present invention relates to the discovery,  
identification, and characterization of novel human  
polynucleotides encoding proteins that share sequence  
similarity with mammalian transporter proteins. The invention  
encompasses the described polynucleotides, host cell expression  
15 systems, the encoded proteins, fusion proteins, polypeptides  
and peptides, antibodies to the encoded proteins and peptides,  
and genetically engineered animals that either lack or  
overexpress the disclosed polynucleotides, antagonists and  
agonists of the proteins, and other compounds that modulate the  
20 expression or activity of the proteins encoded by the disclosed  
polynucleotides, which can be used for diagnosis, drug  
screening, clinical trial monitoring, the treatment of diseases  
and disorders, and cosmetic or nutraceutical applications.

## 2. BACKGROUND OF THE INVENTION

25 Transporter proteins are integral membrane proteins that  
mediate or facilitate the passage of materials across the lipid  
bilayer. Given that the transport of materials across the  
membrane can play an important physiological role, transporter  
proteins are good drug targets. Additionally, one of the  
30 mechanisms of drug resistance involves diseased cells using

WO 02/072774

PCT/US02/07438

cellular transporter systems to export chemotherapeutic agents from the cell. Such mechanisms are particularly relevant to cells manifesting resistance to a multiplicity of drugs.

### 3. SUMMARY OF THE INVENTION

5       The present invention relates to the discovery, identification, and characterization of nucleotides that encode novel human proteins, and the corresponding amino acid sequences of these proteins. The novel human proteins (NHPs), described for the first time herein, share structural  
10 similarity with mammalian sodium/bile co-transporters.

      The novel human nucleic acid sequences described herein, encode alternative proteins/open reading frames (ORFs) of 377 amino acids in length (sodium/bile-like transporter, SEQ ID NO:2), and 438 amino acids in length (sodium/bile-like  
15 transporter, SEQ ID NO:5).

      The invention also encompasses agonists and antagonists of the described NHPs, including small molecules, large molecules, mutant NHPs, or portions thereof, that compete with native NHPs, peptides, and antibodies, as well as nucleotide sequences  
20 that can be used to inhibit the expression of the described NHPs (e.g., antisense and ribozyme molecules, and open reading frame or regulatory sequence replacement constructs) or to enhance the expression of the described NHPs (e.g., expression constructs that place the described polynucleotide under the  
25 control of a strong promoter system), and transgenic animals that express a NHP sequence, or "knock-outs" (which can be conditional) that do not express a functional NHP. Knock-out mice can be produced in several ways, one of which involves the use of mouse embryonic stem cell ("ES cell") lines that contain  
30 gene trap mutations in a murine homolog of at least one of the described NHPs. When the unique NHP sequences described in SEQ

WO 02/072774

PCT/US02/07438

ID NOS:1-6 are "knocked-out" they provide a method of identifying phenotypic expression of the particular gene, as well as a method of assigning function to previously unknown genes. In addition, animals in which the unique NHP sequences described in SEQ ID NOS:1-6 are "knocked-out" provide a unique source in which to elicit antibodies to homologous and orthologous proteins, which would have been previously viewed by the immune system as "self" and therefore would have failed to elicit significant antibody responses.

10        Additionally, the unique NHP sequences described in SEQ ID NOS:1-6 are useful for the identification of protein coding sequences, and mapping a unique gene to a particular chromosome. These sequences identify biologically verified exon splice junctions, as opposed to splice junctions that may have been bioinformatically predicted from genomic sequence alone. The sequences of the present invention are also useful as additional DNA markers for restriction fragment length polymorphism (RFLP) analysis, and in forensic biology.

20        Further, the present invention also relates to processes for identifying compounds that modulate, *i.e.*, act as agonists or antagonists of, NHP expression and/or NHP activity that utilize purified preparations of the described NHPs and/or NHP products, or cells expressing the same. Such compounds can be used as therapeutic agents for the treatment of any of a wide variety of symptoms associated with biological disorders or imbalances.

#### 4. DESCRIPTION OF THE SEQUENCE LISTING AND FIGURES

30        The Sequence Listing provides the sequences of the described NHP ORFs that encode the described NHP amino acid sequences. SEQ ID NOS:3 and 6 describe nucleotides encoding NHP ORFs along with regions of flanking sequence.

WO 02/072774

PCT/US02/07438

## 5. DETAILED DESCRIPTION OF THE INVENTION

The NHPs described for the first time herein are novel proteins that may be expressed in, *inter alia*, human cell lines, brain, pituitary, spinal cord, thymus, spleen, lymph node, trachea, lung, kidney, prostate, testis, thyroid, adrenal gland, pancreas, salivary gland, stomach, small intestine, colon, skeletal muscle, heart, uterus, placenta, mammary gland, adipose, skin, esophagus, bladder, cervix, pericardium, fetal kidney, fetal lung, gall bladder, tongue, aorta, and 6-week old embryos (SEQ ID NOS:1-3), or, *inter alia*, in human cell lines, fetal brain, brain, cerebellum, spinal cord, thymus, spleen, lymph node, bone marrow, trachea, kidney, fetal liver, liver, prostate, testis, thyroid, adrenal gland, pancreas, salivary gland, stomach, small intestine, colon, skeletal muscle, uterus, placenta, mammary gland, esophagus, bladder, cervix, rectum, pericardium, hypothalamus, fetal kidney, fetal lung, gall bladder, tongue, aorta, 6-, 9-, and 12-week old embryos, adenocarcinoma, osteosarcoma, and embryonic carcinoma cells (SEQ ID NOS:4-6).

The present invention encompasses the nucleotides presented in the Sequence Listing, host cells expressing such nucleotides, the expression products of such nucleotides, and: (a) nucleotides that encode mammalian homologs of the described polynucleotides, including the specifically described NHPs, and the NHP products; (b) nucleotides that encode one or more portions of the NHPs that correspond to functional domains, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, the novel regions of any active domain(s); (c) isolated nucleotides that encode mutant versions, engineered or naturally occurring, of the described NHPs in which all or a part of at least one domain is

WO 02/072774

PCT/US02/07438

deleted or altered, and the polypeptide products specified by such nucleotide sequences, including, but not limited to, soluble proteins and peptides in which all or a portion of the signal (or at least one hydrophobic transmembrane) sequence is  
5 deleted; (d) nucleotides that encode chimeric fusion proteins containing all or a portion of a coding region of a NHP, or one of its domains (e.g., a receptor or ligand binding domain, accessory protein/self-association domain, etc.) fused to another peptide or polypeptide; or (e) therapeutic or  
10 diagnostic derivatives of the described polynucleotides, such as oligonucleotides, antisense polynucleotides, ribozymes, dsRNA, or gene therapy constructs, comprising a sequence first disclosed in the Sequence Listing.

As discussed above, the present invention includes the  
15 human DNA sequences presented in the Sequence Listing (and vectors comprising the same), and additionally contemplates any nucleotide sequence encoding a contiguous NHP open reading frame (ORF) that hybridizes to a complement of a DNA sequence presented in the Sequence Listing under highly stringent  
20 conditions, e.g., hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1x SSC/0.1% SDS at 68°C (Ausubel *et al.*, eds., 1989, Current Protocols in Molecular Biology, Vol. I, Green Publishing Associates, Inc., and John Wiley & Sons, Inc., N.Y.,  
25 at p. 2.10.3) and encodes a functionally equivalent expression product. Additionally contemplated are any nucleotide sequences that hybridize to the complement of a DNA sequence that encodes and expresses an amino acid sequence presented in the Sequence Listing under moderately stringent conditions,  
30 e.g., washing in 0.2x SSC/0.1% SDS at 42°C (Ausubel *et al.*, 1989, *supra*), yet still encodes a functionally equivalent NHP product. Functional equivalents of a NHP include naturally

WO 02/072774

PCT/US02/07438

occurring NHPs present in other species, and mutant NHPs, whether naturally occurring or engineered (by site directed mutagenesis, gene shuffling, directed evolution as described in, for example, U.S. Patent No. 5,837,458). The invention  
5 also includes degenerate nucleic acid variants of the disclosed NHP polynucleotide sequences.

Additionally contemplated are polynucleotides encoding NHP ORFs, or their functional equivalents, encoded by polynucleotide sequences that are about 99, 95, 90, or about 85  
10 percent similar or identical to corresponding regions of the nucleotide sequences of the Sequence Listing (as measured by BLAST sequence comparison analysis using, for example, the GCG sequence analysis package, as described herein, using standard default settings).

15 The invention also includes nucleic acid molecules, preferably DNA molecules, that hybridize to, and are therefore the complements of, the described NHP nucleotide sequences. Such hybridization conditions may be highly stringent or less highly stringent, as described herein. In instances where the  
20 nucleic acid molecules are deoxyoligonucleotides ("DNA oligos"), such molecules are generally about 16 to about 100 bases long, or about 20 to about 59 or 80 bases long, or about 34 to about 45 or 47 bases long, or any variation or combination of sizes represented therein that incorporate a  
25 contiguous region of sequence first disclosed in the Sequence Listing. Such oligonucleotides can be used in conjunction with the polymerase chain reaction (PCR) to screen libraries, isolate clones, and prepare cloning and sequencing templates, etc.

30 Alternatively, such NHP oligonucleotides can be used as hybridization probes for screening libraries, and assessing gene expression patterns (particularly using a microarray or

WO 02/072774

PCT/US02/07438

high-throughput "chip" format). Additionally, a series of NHP oligonucleotide sequences, or the complements thereof, can be used to represent all or a portion of the described NHP sequences. An oligonucleotide or polynucleotide sequence first disclosed in at least a portion of one or more of the sequences of SEQ ID NOS:1-6 can be used as a hybridization probe in conjunction with a solid support matrix/substrate (resins, beads, membranes, plastics, polymers, metal or metallized substrates, crystalline or polycrystalline substrates, etc.).

Of particular note are spatially addressable arrays (*i.e.*, gene chips, microtiter plates, etc.) of oligonucleotides and polynucleotides, or corresponding oligopeptides and polypeptides, wherein at least one of the biopolymers present on the spatially addressable array comprises an oligonucleotide or polynucleotide sequence first disclosed in at least one of the sequences of SEQ ID NOS:1-6, or an amino acid sequence encoded thereby. Methods for attaching biopolymers to, or synthesizing biopolymers on, solid support matrices, and conducting binding studies thereon, are disclosed in, *inter alia*, U.S. Patent Nos. 5,700,637, 5,556,752, 5,744,305, 4,631,211, 5,445,934, 5,252,743, 4,713,326, 5,424,186, and 4,689,405, the disclosures of which are herein incorporated by reference in their entirety.

Addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-6 can be used to identify and characterize the temporal and tissue specific expression of a gene. These addressable arrays incorporate oligonucleotide sequences of sufficient length to confer the required specificity, yet be within the limitations of the production technology. The length of these probes is usually within a range of between about 8 to about 2000 nucleotides. Preferably the probes



WO 02/072774

PCT/US02/07438

consist of 60 nucleotides, and more preferably 25 nucleotides, from the sequences first disclosed in SEQ ID NOS:1-6.

For example, a series of oligonucleotide sequences, or the complements thereof, can be used in chip format to represent  
5 all or a portion of the described sequences. The oligonucleotides, typically between about 16 to about 40 (or any whole number within the stated range) nucleotides in length, can partially overlap each other, and/or the sequence may be represented using oligonucleotides that do not overlap.  
10 Accordingly, the described polynucleotide sequences shall typically comprise at least about two or three distinct oligonucleotide sequences of at least about 8 nucleotides in length that are each first disclosed in the described Sequence Listing. Such oligonucleotide sequences can begin at any  
15 nucleotide present within a sequence in the Sequence Listing, and proceed in either a sense (5'-to-3') orientation vis-a-vis the described sequence or in an antisense orientation.

Microarray-based analysis allows the discovery of broad patterns of genetic activity, providing new understanding of  
20 gene functions, and generating novel and unexpected insight into transcriptional processes and biological mechanisms. The use of addressable arrays comprising sequences first disclosed in SEQ ID NOS:1-6 provides detailed information about transcriptional changes involved in a specific pathway,  
25 potentially leading to the identification of novel components, or gene functions that manifest themselves as novel phenotypes.

Probes consisting of sequences first disclosed in SEQ ID NOS:1-6 can also be used in the identification, selection, and validation of novel molecular targets for drug discovery. The  
30 use of these unique sequences permits the direct confirmation of drug targets, and recognition of drug dependent changes in gene expression that are modulated through pathways distinct

WO 02/072774

PCT/US02/07438

from the intended target of the drug. These unique sequences therefore also have utility in defining and monitoring both drug action and toxicity.

As an example of utility, the sequences first disclosed in  
5 SEQ ID NOS:1-6 can be utilized in microarrays, or other assay formats, to screen collections of genetic material from patients who have a particular medical condition. These investigations can also be carried out using the sequences first disclosed in SEQ ID NOS:1-6 *in silico*, and by comparing  
10 previously collected genetic databases and the disclosed sequences using computer software known to those in the art.

Thus the sequences first disclosed in SEQ ID NOS:1-6 can be used to identify mutations associated with a particular disease, and also in diagnostic or prognostic assays.

15 Although the presently described sequences have been specifically described using nucleotide sequence, it should be appreciated that each of the sequences can uniquely be described using any of a wide variety of additional structural attributes, or combinations thereof. For example, a given  
20 sequence can be described by the net composition of the nucleotides present within a given region of the sequence, in conjunction with the presence of one or more specific oligonucleotide sequence(s) first disclosed in SEQ ID NOS:1-6. Alternatively, a restriction map specifying the relative  
25 positions of restriction endonuclease digestion sites, or various palindromic or other specific oligonucleotide sequences, can be used to structurally describe a given sequence. Such restriction maps, which are typically generated by widely available computer programs (e.g., the University of  
30 Wisconsin GCG sequence analysis package, SEQUENCHER 3.0, Gene Codes Corp., Ann Arbor, MI, etc.), can optionally be used in conjunction with one or more discrete nucleotide sequence(s)

WO 02/072774

PCT/US02/07438

present in the sequence that can be described by the relative position of the sequence relative to one or more additional sequence(s) or one or more restriction sites present in the disclosed sequence.

5 For oligonucleotide probes, highly stringent conditions may refer, e.g., to washing in 6x SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligos), 48°C (for 17-base oligos), 55°C (for 20-base oligos), and 60°C (for 23-base oligos). These nucleic acid molecules may encode or act as NHP  
10 antisense molecules, useful, for example, in NHP gene regulation and/or as antisense primers in amplification reactions of NHP nucleic acid sequences. With respect to NHP gene regulation, such techniques can be used to regulate biological functions. Further, such sequences may be used as  
15 part of ribozyme and/or triple helix sequences that are also useful for NHP gene regulation.

Inhibitory antisense or double stranded oligonucleotides can additionally comprise at least one modified base moiety that is selected from the group including, but not limited to,  
20 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine,  
25 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil,  
30 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil,

WO 02/072774

PCT/US02/07438

4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

5           The antisense oligonucleotide can also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

10           In yet another embodiment, the antisense oligonucleotide will comprise at least one modified phosphate backbone selected from the group including, but not limited to, a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

15           In yet another embodiment, the antisense oligonucleotide is an  $\alpha$ -anomeric oligonucleotide. An  $\alpha$ -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the strands run parallel to each other (Gautier *et al.*, 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue *et al.*, 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue *et al.*, 1987, FEBS Lett. 215:327-330). Alternatively, double stranded RNA can be used to disrupt the expression and function of a  
20 targeted NHP.  
25

          Oligonucleotides of the invention can be synthesized by standard methods known in the art, *e.g.*, by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, *etc.*). As examples,  
30 phosphorothioate oligonucleotides can be synthesized by the method of Stein *et al.* (1988, Nucl. Acids Res. 16:3209), and methylphosphonate oligonucleotides can be prepared by use of

WO 02/072774

PCT/US02/07438

controlled pore glass polymer supports (Sarin *et al.*, 1988, Proc. Natl. Acad. Sci. USA 85:7448-7451), *etc.*

Low stringency conditions are well-known to those of skill in the art, and will vary predictably depending on the specific  
5 organisms from which the library and the labeled sequences are derived. For guidance regarding such conditions, see, for example, Sambrook *et al.*, 1989, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (and periodic updates thereof), and Ausubel *et al.*, 1989, *supra*.

10 Alternatively, suitably labeled NHP nucleotide probes can be used to screen a human genomic library using appropriately stringent conditions or by PCR. The identification and characterization of human genomic clones is helpful for identifying polymorphisms (including, but not limited to,  
15 nucleotide repeats, microsatellite alleles, single nucleotide polymorphisms, or coding single nucleotide polymorphisms), determining the genomic structure of a given locus/allele, and designing diagnostic tests. For example, sequences derived from regions adjacent to the intron/exon boundaries of the  
20 human gene can be used to design primers for use in amplification assays to detect mutations within the exons, introns, splice sites (e.g., splice acceptor and/or donor sites), *etc.*, that can be used in diagnostics and pharmacogenomics.

25 For example, the present sequences can be used in restriction fragment length polymorphism (RFLP) analysis to identify specific individuals. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield  
30 unique bands for identification (as generally described in U.S. Patent No. 5,272,057, incorporated herein by reference). In addition, the sequences of the present invention can be used to

WO 02/072774

PCT/US02/07438

provide polynucleotide reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (*i.e.*,  
5 another DNA sequence that is unique to a particular individual). Actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme generated fragments.

Further, a NHP gene homolog can be isolated from nucleic  
10 acid from an organism of interest by performing PCR using two degenerate or "wobble" oligonucleotide primer pools designed on the basis of amino acid sequences within the NHP products disclosed herein. The template for the reaction may be total RNA, mRNA, and/or cDNA obtained by reverse transcription of  
15 mRNA, prepared from human or non-human cell lines or tissue known to express, or suspected of expressing, an allele of a NHP gene.

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequence of the  
20 desired NHP gene. The PCR fragment can then be used to isolate a full length cDNA clone by a variety of methods. For example, the amplified fragment can be labeled and used to screen a cDNA library, such as a bacteriophage cDNA library. Alternatively, the labeled fragment can be used to isolate genomic clones via  
25 the screening of a genomic library.

PCR technology can also be used to isolate full length cDNA sequences. For example, RNA can be isolated, following standard procedures, from an appropriate cellular or tissue source (*i.e.*, one known to express, or suspected of expressing,  
30 a NHP gene). A reverse transcription (RT) reaction can be performed on the RNA using an oligonucleotide primer specific for the most 5' end of the amplified fragment for the priming

WO 02/072774

PCT/US02/07438

of first strand synthesis. The resulting RNA/DNA hybrid may then be "tailed" using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a complementary  
5 primer. Thus, cDNA sequences upstream of the amplified fragment can be isolated. For a review of cloning strategies that can be used, see, e.g., Sambrook et al., 1989, *supra*.

A cDNA encoding a mutant NHP sequence can be isolated, for example, by using PCR. In this case, the first cDNA strand may  
10 be synthesized by hybridizing an oligo-dT oligonucleotide to mRNA isolated from tissue known to express, or suspected of expressing, a NHP, in an individual putatively carrying a mutant NHP allele, and by extending the new strand with reverse transcriptase. The second strand of the cDNA is then  
15 synthesized using an oligonucleotide that hybridizes specifically to the 5' end of the normal sequence. Using these two primers, the product is then amplified via PCR, optionally cloned into a suitable vector, and subjected to DNA sequence analysis through methods well-known to those of skill in the  
20 art. By comparing the DNA sequence of the mutant NHP allele to that of a corresponding normal NHP allele, the mutation(s) responsible for the loss or alteration of function of the mutant NHP gene product can be ascertained.

Alternatively, a genomic library can be constructed using  
25 DNA obtained from an individual suspected of carrying, or known to carry, a mutant NHP allele (e.g., a person manifesting a NHP-associated phenotype such as, for example, obesity, high blood pressure, connective tissue disorders, infertility, etc.), or a cDNA library can be constructed using RNA from a  
30 tissue known to express, or suspected of expressing, a mutant NHP allele. A normal NHP gene, or any suitable fragment thereof, can then be labeled and used as a probe to identify

WO 02/072774

PCT/US02/07438

the corresponding mutant NHP allele in such libraries. Clones containing mutant NHP sequences can then be purified and subjected to sequence analysis according to methods well-known to those skilled in the art.

5        Additionally, an expression library can be constructed utilizing cDNA synthesized from, for example, RNA isolated from a tissue known to express, or suspected of expressing, a mutant NHP allele in an individual suspected of carrying, or known to carry, such a mutant allele. In this manner, gene products  
10        made by the putatively mutant tissue can be expressed and screened using standard antibody screening techniques in conjunction with antibodies raised against a normal NHP product, as described below (for screening techniques, see, for example, Harlow and Lane, eds., 1988, "Antibodies: A Laboratory  
15        Manual", Cold Spring Harbor Press, Cold Spring Harbor, N.Y.).

         Additionally, screening can be accomplished by screening with labeled NHP fusion proteins, such as, for example, alkaline phosphatase-NHP or NHP-alkaline phosphatase fusion proteins. In cases where a NHP mutation results in an  
20        expression product with altered function (e.g., as a result of a missense or a frameshift mutation), polyclonal antibodies to a NHP are likely to cross-react with a corresponding mutant NHP expression product. Library clones detected via their reaction with such labeled antibodies can be purified and subjected to  
25        sequence analysis according to methods well-known in the art.

         The invention also encompasses: (a) DNA vectors that contain any of the foregoing NHP coding sequences and/or their complements (*i.e.*, antisense); (b) DNA expression vectors that contain any of the foregoing NHP coding sequences operatively  
30        associated with a regulatory element that directs the expression of the coding sequences (for example, baculovirus as described in U.S. Patent No. 5,869,336 herein incorporated by



WO 02/072774

PCT/US02/07438

reference); (c) genetically engineered host cells that contain any of the foregoing NHP coding sequences operatively associated with a regulatory element that directs the expression of the coding sequences in the host cell; and (d)  
5 genetically engineered host cells that express an endogenous NHP sequence under the control of an exogenously introduced regulatory element (*i.e.*, gene activation). As used herein, regulatory elements include, but are not limited to, inducible and non-inducible promoters, enhancers, operators, and other  
10 elements known to those skilled in the art that drive and regulate expression. Such regulatory elements include, but are not limited to, the cytomegalovirus (hCMV) immediate early gene, regulatable, viral elements (particularly retroviral LTR promoters), the early or late promoters of SV40 or adenovirus,  
15 the *lac* system, the *trp* system, the *TAC* system, the *TRC* system, the major operator and promoter regions of phage lambda, the control regions of fd coat protein, the promoter for 3-phosphoglycerate kinase (PGK), the promoters of acid phosphatase, and the promoters of the yeast  $\alpha$ -mating factors.

20 The present invention also encompasses antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists and agonists of a NHP, as well as compounds or nucleotide constructs that inhibit expression of a NHP sequence (transcription factor inhibitors, antisense and ribozyme  
25 molecules, or open reading frame sequence or regulatory sequence replacement constructs), or promote the expression of a NHP (*e.g.*, expression constructs in which NHP coding sequences are operatively associated with expression control elements such as promoters, promoter/enhancers, *etc.*).

30 The NHPs or NHP peptides, NHP fusion proteins, NHP nucleotide sequences, antibodies, antagonists and agonists can be useful for the detection of mutant NHPs, or inappropriately

WO 02/072774

PCT/US02/07438

expressed NHPs, for the diagnosis of disease. The NHP proteins or peptides, NHP fusion proteins, NHP nucleotide sequences, host cell expression systems, antibodies, antagonists, agonists and genetically engineered cells and animals can be used for  
5 screening for drugs (or high throughput screening of combinatorial libraries) effective in the treatment of the symptomatic or phenotypic manifestations of perturbing the normal function of a NHP in the body. The use of engineered host cells and/or animals may offer an advantage in that such  
10 systems allow not only for the identification of compounds that bind to the endogenous receptor for a NHP, but can also identify compounds that trigger NHP-mediated activities or pathways.

Finally, the NHP products can be used as therapeutics.  
15 For example, soluble derivatives such as NHP peptides/domains corresponding to NHPs, NHP fusion protein products (especially NHP-Ig fusion proteins, *i.e.*, fusions of a NHP, or a domain of a NHP, to an IgFc), NHP antibodies and anti-idiotypic antibodies (including Fab fragments), antagonists or agonists  
20 (including compounds that modulate or act on downstream targets in a NHP-mediated pathway) can be used to directly treat diseases or disorders.. For instance, the administration of an effective amount of a soluble NHP, a NHP-IgFc fusion protein, or an anti-idiotypic antibody (or its Fab) that mimics a NHP  
25 could activate or effectively antagonize the endogenous NHP receptor. Nucleotide constructs encoding such NHP products can be used to genetically engineer host cells to express such products *in vivo*; these genetically engineered cells function as "bioreactors" in the body delivering a continuous supply of  
30 a NHP, a NHP peptide, or a NHP fusion protein to the body. Nucleotide constructs encoding functional NHPs, mutant NHPs, as well as antisense and ribozyme molecules can also be used in

WO 02/072774

PCT/US02/07438

"gene therapy" approaches for the modulation of NHP expression. Thus, the invention also encompasses pharmaceutical formulations and methods for treating biological disorders.

Various aspects of the invention are described in greater  
5 detail in the subsections below.

### 5.1 THE NHP SEQUENCES

The cDNA sequences and the corresponding deduced amino acid sequences of the described NHPs are presented in the  
10 Sequence Listing. The NHP nucleotides were obtained from clustered human genomic sequences, testis and mammary transcript RACE products, ESTs, and human cDNAs made from fetal kidney, lymph node, trachea, lung, mammary gland, bone marrow, and pericardium (SEQ ID NOS:1-3), and kidney, fetal kidney,  
15 mammary gland, and prostate (SEQ ID NOS:4-6) mRNAs (Edge Biosystems, Gaithersburg, MD, and Clontech, Palo Alto, CA).

One polymorphisms was found, a C/T polymorphism at the nucleotide position corresponding to nucleotide 426 of, for example, SEQ ID NO:4, both of which result in an asn being  
20 present at corresponding amino acid position 142 of SEQ ID NO:5. SEQ ID NOS:1-3 are apparently encoded on human chromosome 4 (see GENBANK accession no. AC079237), and SEQ ID NOS:4-6 are apparently encoded on human chromosome 8 (see GENBANK accession no. AC024371). Accordingly, the described  
25 sequences are useful for mapping and/or defining the corresponding coding regions of the human genome.

An additional application of the described novel human polynucleotide sequences is their use in the molecular mutagenesis/evolution of proteins that are at least partially  
30 encoded by the described novel sequences using, for example, polynucleotide shuffling or related methodologies. Such approaches are described in U.S. Patent Nos. 5,830,721 and

WO 02/072774

PCT/US02/07438

5,837,458, which are herein incorporated by reference in their entirety.

NHP gene products can also be expressed in transgenic animals. Animals of any species, including, but not limited  
5 to, worms, mice, rats, rabbits, guinea pigs, pigs, micro-pigs, birds, goats, and non-human primates, e.g., baboons, monkeys, and chimpanzees, may be used to generate NHP transgenic animals.

Any technique known in the art may be used to introduce a  
10 NHP transgene into animals to produce the founder lines of transgenic animals. Such techniques include, but are not limited to, pronuclear microinjection (Hoppe and Wagner, 1989, U.S. Patent No. 4,873,191); retrovirus-mediated gene transfer into germ lines (Van der Putten *et al.*, 1985, Proc. Natl. Acad.  
15 Sci. USA 82:6148-6152); gene targeting in embryonic stem cells (Thompson *et al.*, 1989, Cell 56:313-321); electroporation of embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814); and sperm-mediated gene transfer (Lavitrano *et al.*, 1989, Cell 57:717-723); etc. For a review of such techniques, see Gordon, 1989,  
20 Transgenic Animals, Intl. Rev. Cytol. 115:171-229, which is incorporated by reference herein in its entirety.

The present invention provides for transgenic animals that carry a NHP transgene in all their cells, as well as animals that carry a transgene in some, but not all their cells, i.e.,  
25 mosaic animals or somatic cell transgenic animals. A transgene may be integrated as a single transgene, or in concatamers, e.g., head-to-head tandems or head-to-tail tandems. A transgene may also be selectively introduced into and activated in a particular cell-type by following, for example, the  
30 teaching of Lasko *et al.*, 1992, Proc. Natl. Acad. Sci. USA 89:6232-6236. The regulatory sequences required for such a cell-type specific activation will depend upon the particular

WO 02/072774

PCT/US02/07438

cell-type of interest, and will be apparent to those of skill in the art.

When it is desired that a NHP transgene be integrated into the chromosomal site of the endogenous NHP gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous NHP gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous NHP gene (*i.e.*, "knockout" animals).

The transgene can also be selectively introduced into a particular cell-type, thus inactivating the endogenous NHP gene in only that cell-type, by following, for example, the teaching of Gu *et al.*, 1994, Science 265:103-106. The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell-type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant NHP gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to assay whether integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques that include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and RT-PCR. Samples of NHP gene-expressing tissue may also be evaluated immunocytochemically using antibodies specific for the NHP transgene product.

The present invention also provides for "knock-in" animals. Knock-in animals are those in which a polynucleotide

WO 02/072774

PCT/US02/07438

sequence (*i.e.*, a gene or a cDNA) that the animal does not naturally have in its genome is inserted in such a way that it is expressed. Examples include, but are not limited to, a human gene or cDNA used to replace its murine ortholog in the mouse, a murine cDNA used to replace the murine gene in the mouse, and a human gene or cDNA or murine cDNA that is tagged with a reporter construct used to replace the murine ortholog or gene in the mouse. Such replacements can occur at the locus of the murine ortholog or gene, or at another specific site. Such knock-in animals are useful for the *in vivo* study, testing and validation of, *intra alia*, human drug targets, as well as for compounds that are directed at the same and therapeutic proteins.

## 5.2 NHPS AND NHP POLYPEPTIDES

NHPS, NHP polypeptides, NHP peptide fragments, mutated, truncated, or deleted forms of the NHPS, and/or NHP fusion proteins can be prepared for a variety of uses. These uses include, but are not limited to, the generation of antibodies, as reagents in diagnostic assays, for the identification of other cellular gene products related to a NHP, and as reagents in assays for screening for compounds that can be used as pharmaceutical reagents useful in the therapeutic treatment of mental, biological, or medical disorders and diseases. Given the similarity information and expression data, the described NHPS can be targeted (by drugs, oligos, antibodies, etc.) in order to treat disease, or to therapeutically augment the efficacy of, for example, chemotherapeutic agents used in the treatment of breast or prostate cancer.

The Sequence Listing discloses the amino acid sequences encoded by the described NHP polynucleotides. The NHPS typically display initiator methionines in DNA sequence

WO 02/072774

PCT/US02/07438

contexts consistent with a translation initiation site. SEQ ID NO:5 displays a signal type sequence similar to those often found on membrane proteins; however, all of the described proteins display multiple transmembrane hydrophobic domains  
5 typical of membrane associated proteins.

The NHP amino acid sequences of the invention include the amino acid sequence presented in the Sequence Listing, as well as analogues and derivatives thereof. Further, corresponding NHP homologues from other species are encompassed by the  
10 invention. In fact, any NHP protein encoded by the NHP nucleotide sequences described herein are within the scope of the invention, as are any novel polynucleotide sequences encoding all or any novel portion of an amino acid sequence presented in the Sequence Listing. The degenerate nature of  
15 the genetic code is well-known, and, accordingly, each amino acid presented in the Sequence Listing is generically representative of the well-known nucleic acid "triplet" codon, or in many cases codons, that can encode the amino acid. As such, as contemplated herein, the amino acid sequences  
20 presented in the Sequence Listing, when taken together with the genetic code (see, for example, Table 4-1 at page 109 of "Molecular Cell Biology", 1986, J. Darnell et al., eds., Scientific American Books, New York, N.Y., herein incorporated by reference), are generically representative of all the  
25 various permutations and combinations of nucleic acid sequences that can encode such amino acid sequences.

The invention also encompasses proteins that are functionally equivalent to the NHPs encoded by the presently described nucleotide sequences, as judged by any of a number of  
30 criteria, including, but not limited to, the ability to bind and cleave a substrate of a NHP, the ability to effect an identical or complementary downstream pathway, or a change in

WO 02/072774

PCT/US02/07438

cellular metabolism (e.g., proteolytic activity, ion flux, tyrosine phosphorylation, transport, etc.). Such functionally equivalent NHP proteins include, but are not limited to, additions or substitutions of amino acid residues within the amino acid sequence encoded by the NHP nucleotide sequences described herein, but that result in a silent change, thus producing a functionally equivalent expression product. Amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid.

A variety of host-expression vector systems can be used to express the NHP nucleotide sequences of the invention. Where, as in the present instance, the NHP peptides or polypeptides are thought to be from a membrane protein, the hydrophobic regions of the protein can be excised and the resulting soluble peptides or polypeptides can be recovered from the culture media. Such expression systems also encompass engineered host cells that express a NHP, or a functional equivalent, *in situ*. Purification or enrichment of a NHP from such expression systems can be accomplished using appropriate detergents and lipid micelles and methods well-known to those skilled in the art. However, such engineered host cells themselves may be used in situations where it is important not only to retain the structural and functional characteristics of a NHP, but to



WO 02/072774

PCT/US02/07438

assess biological activity, e.g., in certain drug screening assays.

The expression systems that may be used for purposes of the invention include, but are not limited to, microorganisms  
5 such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing NHP nucleotide sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing NHP nucleotide sequences;  
10 insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing NHP nucleotide sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant  
15 plasmid expression vectors (e.g., Ti plasmid) containing NHP nucleotide sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3) harboring recombinant expression constructs containing NHP nucleotide sequences and promoters derived from the genome of mammalian cells (e.g., metallothionein promoter)  
20 or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

In bacterial systems, a number of expression vectors may be advantageously selected depending upon the use intended for the NHP product being expressed. For example, when a large  
25 quantity of such a protein is to be produced for the generation of pharmaceutical compositions of or containing a NHP, or for raising antibodies to a NHP, vectors that direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not  
30 limited to, the *E. coli* expression vector pUR278 (Ruther et al., 1983, EMBO J. 2:1791), in which a NHP coding sequence may be ligated individually into the vector in-frame with the *lacZ*

WO 02/072774

PCT/US02/07438

coding region so that a fusion protein is produced; pIN vectors (Inouye and Inouye, 1985, Nucleic Acids Res. 13:3101-3109; Van Heeke and Schuster, 1989, J. Biol. Chem. 264:5503-5509); and the like. pGEX vectors (Pharmacia or American Type Culture  
5 Collection) can also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione.  
10 The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target expression product can be released from the GST moiety.

In an exemplary insect system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to  
15 express foreign polynucleotide sequences. The virus grows in *Spodoptera frugiperda* cells. A NHP coding sequence can be cloned individually into a non-essential region (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).  
20 Successful insertion of a NHP coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera*  
25 *frugiperda* cells in which the inserted sequence is expressed (e.g., see Smith et al., 1983, J. Virol. 46:584; Smith, U.S. Patent No. 4,215,051).

In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an  
30 adenovirus is used as an expression vector, the NHP nucleotide sequence of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late

WO 02/072774

PCT/US02/07438

promoter and tripartite leader sequence. This chimeric sequence may then be inserted in the adenovirus genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a NHP product in infected hosts (e.g., see Logan and Shenk, 1984, Proc. Natl. Acad. Sci. USA 81:3655-3659). Specific initiation signals may also be required for efficient translation of inserted NHP nucleotide sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire NHP gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. However, in cases where only a portion of a NHP coding sequence is inserted, exogenous translational control signals, including, perhaps, the ATG initiation codon, may be provided. Furthermore, the initiation codon should be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bitter et al., 1987, Methods in Enzymol. 153:516-544).

In addition, a host cell strain may be chosen that modulates the expression of the inserted sequences, or modifies and processes the expression product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-

WO 02/072774

PCT/US02/07438

translational processing and modification of proteins and expression products. Appropriate cell lines or host systems can be chosen to ensure the desired modification and processing of the foreign protein expressed. To this end, eukaryotic host  
5 cells that possess the cellular machinery for the desired processing of the primary transcript, glycosylation, and phosphorylation of the expression product may be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and in particular,  
10 human cell lines.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express the NHP sequences described herein can be engineered. Rather than using expression vectors that  
15 contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign  
20 DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in  
25 turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines that express the NHP product. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that affect the endogenous activity of the NHP product.

30 A number of selection systems may be used, including, but not limited to, the herpes simplex virus thymidine kinase (Wigler et al., 1977, Cell 11:223), hypoxanthine-guanine

WO 02/072774

PCT/US02/07438

phosphoribosyltransferase (Szybalska and Szybalski, 1962, Proc. Natl. Acad. Sci. USA 48:2026), and adenine phosphoribosyltransferase (Lowy et al., 1980, Cell 22:817) genes, which can be employed in tk<sup>-</sup>, hgp<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77:3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78:1527); gpt, which confers resistance to mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78:2072); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150:1); and hyg<sup>r</sup>, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30:147).

Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. Another exemplary system allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., 1991, Proc. Natl. Acad. Sci. USA 88:8972-8976). In this system, the sequence of interest is subcloned into a vaccinia recombination plasmid such that the sequence's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni<sup>2+</sup>-nitriloacetic acid-agarose columns, and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

Also encompassed by the present invention are fusion proteins that direct a NHP to a target organ and/or facilitate transport across the membrane into the cytosol. Conjugation of NHPs to antibody molecules or their Fab fragments could be used to target cells bearing a particular epitope. Attaching an

WO 02/072774

PCT/US02/07438

appropriate signal sequence to a NHP would also transport a NHP to a desired location within the cell. Alternatively targeting of a NHP or its nucleic acid sequence might be achieved using liposome or lipid complex based delivery systems. Such technologies are described in "Liposomes: A Practical Approach", New, R.R.C., ed., Oxford University Press, N.Y., and in U.S. Patent Nos. 4,594,595, 5,459,127, 5,948,767 and 6,110,490 and their respective disclosures, which are herein incorporated by reference in their entirety. Additionally embodied are novel protein constructs engineered in such a way that they facilitate transport of NHPs to a target site or desired organ, where they cross the cell membrane and/or the nucleus where the NHPs can exert their functional activity. This goal may be achieved by coupling of a NHP to a cytokine or other ligand that provides targeting specificity, and/or to a protein transducing domain (see generally U.S. Provisional Patent Application Ser. Nos. 60/111,701 and 60/056,713, both of which are herein incorporated by reference, for examples of such transducing sequences), to facilitate passage across cellular membranes, and can optionally be engineered to include nuclear localization signals.

### 5.3 ANTIBODIES TO NHP PRODUCTS

Antibodies that specifically recognize one or more epitopes of a NHP, epitopes of conserved variants of a NHP, or peptide fragments of a NHP, are also encompassed by the invention. Such antibodies include, but are not limited to, polyclonal antibodies, monoclonal antibodies (mAbs), humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')<sub>2</sub> fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, and epitope-binding fragments of any of the above.

WO 02/072774

PCT/US02/07438

The antibodies of the invention may be used, for example, in the detection of a NHP in a biological sample and may, therefore, be utilized as part of a diagnostic or prognostic technique whereby patients may be tested for abnormal amounts of a NHP. Such antibodies may also be utilized in conjunction with, for example, compound screening schemes for the evaluation of the effect of test compounds on expression and/or activity of a NHP expression product. Additionally, such antibodies can be used in conjunction with gene therapy to, for example, evaluate normal and/or engineered NHP-expressing cells prior to their introduction into a patient. Such antibodies may additionally be used in methods for the inhibition of abnormal NHP activity. Thus, such antibodies may be utilized as a part of treatment methods.

For the production of antibodies, various host animals may be immunized by injection with a NHP, a NHP peptide (e.g., one corresponding to a functional domain of a NHP), truncated NHP polypeptides (NHP in which one or more domains have been deleted), functional equivalents of the NHP, or mutated variants of the NHP. Such host animals may include, but are not limited to, pigs, rabbits, mice, goats, and rats, to name but a few. Various adjuvants may be used to increase the immunological response, depending on the host species, including, but not limited to, Freund's adjuvant (complete and incomplete), mineral salts such as aluminum hydroxide or aluminum phosphate, chitosan, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Alternatively, the immune response could be enhanced by combination and/or coupling with molecules such as keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid,

WO 02/072774

PCT/US02/07438

ovalbumin, cholera toxin, or fragments thereof. Polyclonal antibodies are heterogeneous populations of antibody molecules derived from the sera of the immunized animals.

Monoclonal antibodies, which are homogeneous populations  
5 of antibodies to a particular antigen, can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique of Kohler and Milstein, (1975, *Nature* 256:495-497; and U.S. Patent No.  
10 4,376,110), the human B-cell hybridoma technique (Kosbor *et al.*, 1983, *Immunology Today* 4:72; Cole *et al.*, 1983, *Proc. Natl. Acad. Sci. USA* 80:2026-2030), and the EBV-hybridoma technique (Cole *et al.*, 1985, *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96). Such antibodies may  
15 be of any immunoglobulin class, including IgG, IgM, IgE, IgA, and IgD, and any subclass thereof. The hybridomas producing the mAbs of this invention may be cultivated *in vitro* or *in vivo*. Production of high titers of mAbs *in vivo* makes this the presently preferred method of production.

20 In addition, techniques developed for the production of "chimeric antibodies" (Morrison *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger *et al.*, 1984, *Nature*, 312:604-608; Takeda *et al.*, 1985, *Nature*, 314:452-454), by splicing the genes from a mouse antibody molecule of appropriate antigen  
25 specificity together with genes from a human antibody molecule of appropriate biological activity, can be used. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin  
30 constant region. Such technologies are described in U.S. Patent Nos. 6,114,598, 6,075,181 and 5,877,397 and their respective disclosures, which are herein incorporated by



WO 02/072774

PCT/US02/07438

reference in their entirety. Also encompassed by the present invention is the use of fully humanized monoclonal antibodies, as described in U.S. Patent No. 6,150,584 and respective disclosures, which are herein incorporated by reference in  
5 their entirety.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, 1988, *Science* 242:423-426; Huston et al., 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Ward et al., 1989, *Nature* 341:544-  
10 546) can be adapted to produce single chain antibodies against NHP expression products. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

15 Antibody fragments that recognize specific epitopes may be generated by known techniques. For example, such fragments include, but are not limited to: F(ab')<sub>2</sub> fragments, which can be produced by pepsin digestion of an antibody molecule; and Fab fragments, which can be generated by reducing the disulfide  
20 bridges of F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed (Huse et al., 1989, *Science*, 246:1275-1281) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

Antibodies to a NHP can, in turn, be utilized to generate  
25 anti-idiotypic antibodies that "mimic" a given NHP, using techniques well-known to those skilled in the art (see, e.g., Greenspan and Bona, 1993, *FASEB J.* 7:437-444; and Nissinoff, 1991, *J. Immunol.* 147:2429-2438). For example, antibodies that  
30 bind to a NHP domain and competitively inhibit the binding of the NHP to its cognate receptor can be used to generate anti-idiotypes that "mimic" the NHP and, therefore, bind and activate or neutralize a receptor. Such anti-idiotypic

WO 02/072774

PCT/US02/07438

antibodies, or Fab fragments of such anti-idiotypes, can be used in therapeutic regimens involving a NHP-mediated pathway.

Additionally, given the high degree of relatedness of mammalian NHPs, NHP knock-out mice (having never seen a NHP, and thus never been tolerized to a NHP) have a unique utility, as they can be advantageously applied to the generation of antibodies against the disclosed mammalian NHPs (*i.e.*, a NHP will be immunogenic in NHP knock-out animals).

The present invention is not to be limited in scope by the specific embodiments described herein, which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein, will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims. All cited publications, patents, and patent applications are herein incorporated by reference in their entirety.

WO 02/072774

PCT/US02/07438

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising at least 59 contiguous nucleotides from SEQ ID NO:1.

5

2. An isolated nucleic acid molecule comprising a nucleotide sequence that:

(a) encodes the amino acid sequence shown in SEQ ID NO:2; and

10 (b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO:1 or the complement thereof.

3. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO:2.

4. A substantially isolated protein having the transporter activity of the protein shown in SEQ ID NO:2, which is encoded by a nucleotide sequence that hybridizes to SEQ ID NO:1 under highly stringent conditions.

5. A substantially isolated protein according to Claim 4 comprising the amino acid sequence of SEQ ID NO:2.

25

6. An isolated nucleic acid molecule comprising at least 47 contiguous nucleotides from SEQ ID NO:4.

7. An isolated nucleic acid molecule comprising a nucleotide sequence that:

30 (a) encodes the amino acid sequence shown in SEQ ID NO:5; and

WO 02/072774

PCT/US02/07438

(b) hybridizes under stringent conditions to the nucleotide sequence of SEQ ID NO:4 or the complement thereof.

5                   8.    An isolated nucleic acid molecule comprising a nucleotide sequence that encodes the amino acid sequence shown in SEQ ID NO:4.

                  9.    A substantially isolated protein having the  
10   transporter activity of the protein shown in SEQ ID NO:5, which is encoded by a nucleotide sequence that hybridizes to SEQ ID NO:4 under highly stringent conditions.

                  10.   A substantially isolated protein according to  
15   Claim 9 comprising the amino acid sequence of SEQ ID NO:5.

WO 02/072774

PCT/US02/07438

SEQUENCE LISTING

<110> LEXICON GENETICS INCORPORATED

<120> Novel Human Transporter Proteins and Polynucleotides Encoding the Same

<130> LEX-0314-PCT

<150> US 60/275,009

<151> 2001-03-12

<150> US 60/284,152

<151> 2001-04-17

<160> 6

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1134

<212> DNA

<213> Homo sapiens

<400> 1

atgagagcca	attgttccag	cagctcagcc	tgccctgcc	acagttcaga	ggaggagctg	60
ccagtgggac	tggaggtgca	tggaaacctg	gagctcgttt	tcacagtggg	gtccactgtg	120
atgatggggc	tgctcatgtt	ctctttggga	tggtccgtgg	agatccggaa	gctgtggctg	180
cacatcagga	gaccctgggg	cattgctgtg	ggactgctct	gccagtttgg	gctcatgcct	240
tttacagctt	atctcctggc	cattagcttt	tctctgaagc	cagtccaagc	tattgctgtt	300
ctcatcatgg	gctgctgccc	ggggggcacc	atctctaaca	ttttcacctt	ctgggttgat	360
ggagatatgg	atctcagcat	cagtatgaca	acctgttcca	ccgtggccgc	cctgggaatg	420
atgccactct	gcatttatct	ctacacctgg	tcctggagtc	ttcagcagaa	tctcaccatt	480
ccttatcaga	acataggaat	tacccttggt	tgccctgacca	ttcctgtggc	ctttggtgtc	540
tatgtgaatt	acagatggcc	aaaacaatcc	aaaatcattc	tcaagattgg	ggccgttggt	600
ggtgggggtc	tccttctggt	ggtcgcagtt	gctggtgttg	tcctggcgaa	aggatcttgg	660
aattcagaca	tcacccttct	gaccatcagt	ttcatcttcc	ctttgattgg	ccatgtcacg	720
ggttttctgc	tggcactttt	taccaccag	tcctggcaaa	ggtgcaggac	aatttcctta	780
gaaactggag	ctcagaatat	tcagatgtgc	atcaccatgc	tcagttatc	tttactgct	840
gagcacttgg	tccagatgtt	gagtttccca	ctggcctatg	gactcttcca	gctgatagat	900
ggattttctta	ttgttgcagc	atatcagacg	tacaagagga	gattgaagaa	caaacatgga	960
aaaaagaact	caggttgcac	agaagtctgc	catacgagga	aatcgacttc	ttccagagag	1020
accaatgcct	tccttgagggt	gaatgaagaa	ggtgccatca	ctcctggggc	accagggcca	1080
atggattgcc	acagggctct	cgagccagtt	ggccacatca	cttcatgtga	atag	1134

<210> 2

<211> 377

<212> PRT

<213> Homo sapiens

<400> 2

Met	Arg	Ala	Asn	Cys	Ser	Ser	Ser	Ser	Ala	Cys	Pro	Ala	Asn	Ser	Ser
1				5					10					15	
Glu	Glu	Glu	Leu	Pro	Val	Gly	Leu	Glu	Val	His	Gly	Asn	Leu	Glu	Leu

WO 02/072774

PCT/US02/07438

Val	Phe	Thr	Val	Val	Ser	Thr	Val	Met	Met	Gly	Leu	Leu	Met	Phe	Ser
	35						40					45			
Leu	Gly	Cys	Ser	Val	Glu	Ile	Arg	Lys	Leu	Trp	Ser	His	Ile	Arg	Arg
	50					55					60				
Pro	Trp	Gly	Ile	Ala	Val	Gly	Leu	Leu	Cys	Gln	Phe	Gly	Leu	Met	Pro
65					70					75					80
Phe	Thr	Ala	Tyr	Leu	Leu	Ala	Ile	Ser	Phe	Ser	Leu	Lys	Pro	Val	Gln
				85					90					95	
Ala	Ile	Ala	Val	Leu	Ile	Met	Gly	Cys	Cys	Pro	Gly	Gly	Thr	Ile	Ser
			100					105					110		
Asn	Ile	Phe	Thr	Phe	Trp	Val	Asp	Gly	Asp	Met	Asp	Leu	Ser	Ile	Ser
	115						120					125			
Met	Thr	Thr	Cys	Ser	Thr	Val	Ala	Ala	Leu	Gly	Met	Met	Pro	Leu	Cys
	130					135					140				
Ile	Tyr	Leu	Tyr	Thr	Trp	Ser	Trp	Ser	Leu	Gln	Gln	Asn	Leu	Thr	Ile
145					150					155					160
Pro	Tyr	Gln	Asn	Ile	Gly	Ile	Thr	Leu	Val	Cys	Leu	Thr	Ile	Pro	Val
			165						170					175	
Ala	Phe	Gly	Val	Tyr	Val	Asn	Tyr	Arg	Trp	Pro	Lys	Gln	Ser	Lys	Ile
			180					185					190		
Ile	Leu	Lys	Ile	Gly	Ala	Val	Val	Gly	Gly	Val	Leu	Leu	Leu	Val	Val
	195						200					205			
Ala	Val	Ala	Gly	Val	Val	Leu	Ala	Lys	Gly	Ser	Trp	Asn	Ser	Asp	Ile
	210					215					220				
Thr	Leu	Leu	Thr	Ile	Ser	Phe	Ile	Phe	Pro	Leu	Ile	Gly	His	Val	Thr
225					230					235					240
Gly	Phe	Leu	Leu	Ala	Leu	Phe	Thr	His	Gln	Ser	Trp	Gln	Arg	Cys	Arg
				245					250					255	
Thr	Ile	Ser	Leu	Glu	Thr	Gly	Ala	Gln	Asn	Ile	Gln	Met	Cys	Ile	Thr
			260					265					270		
Met	Leu	Gln	Leu	Ser	Phe	Thr	Ala	Glu	His	Leu	Val	Gln	Met	Leu	Ser
	275						280					285			
Phe	Pro	Leu	Ala	Tyr	Gly	Leu	Phe	Gln	Leu	Ile	Asp	Gly	Phe	Leu	Ile
	290					295					300				
Val	Ala	Ala	Tyr	Gln	Thr	Tyr	Lys	Arg	Arg	Leu	Lys	Asn	Lys	His	Gly
305					310					315					320
Lys	Lys	Asn	Ser	Gly	Cys	Thr	Glu	Val	Cys	His	Thr	Arg	Lys	Ser	Thr
				325					330					335	
Ser	Ser	Arg	Glu	Thr	Asn	Ala	Phe	Leu	Glu	Val	Asn	Glu	Glu	Gly	Ala
			340					345					350		
Ile	Thr	Pro	Gly	Pro	Pro	Gly	Pro	Met	Asp	Cys	His	Arg	Ala	Leu	Glu
	355						360					365			
Pro	Val	Gly	His	Ile	Thr	Ser	Cys	Glu							
	370					375									

<210> 3  
 <211> 1600  
 <212> DNA  
 <213> Homo sapiens

<400> 3  
 gtgataatta cttttataat gccacttgtg aaaaaattga tcagattagg atgaatcacc 60  
 ttgctggcca acagttattg gaatgattct ccatgtgtga cttcgttgca ctattacaaa 120  
 atgtggcagg atagacctgc ccagccattg ttgccgatgt tcatttgtaa tgctgcctta 180  
 aggagatgag gagatgagag ccaattgttc cagcagctca gcctgccctg ccaacagttc 240

WO 02/072774

PCT/US02/07438

agaggaggag	ctgccagtg	gactggagg	gcatggaaac	ctggagctcg	ttttcacagt	300
gggtgtccact	gtgatgatgg	ggctgtctcat	gttctctttg	ggatgttccg	tggagatccg	360
gaagctgtgg	tcgcacatca	ggagaccctg	gggcattgct	gtgggactgc	tctgccagtt	420
tgggctcatg	cctttttacag	cttatctcct	ggccattagc	ttttctctga	agccagtcca	480
agctattgct	gttctcatca	tgggctgctg	cccggggggc	accatctcta	acattttcac	540
cttctgggtt	gatggagata	tggatctcag	catcagtatg	acaacctgtt	ccaccgtggc	600
cgccctggga	atgatgccac	tctgcattta	tctctacacc	tggctctgga	gtcttcagca	660
gaatctcacc	attccttatc	agaacatagg	aattaccctt	gtgtgcctga	ccattcctgt	720
ggccttttgt	gtctatgtga	attacagatg	gccaaaacaa	tccaaaatca	ttctcaagat	780
tggggccggt	gttggtgggg	tcctccttct	gggtgctgca	gttgctgggt	tggctcctggc	840
gaaaggatct	tgggaattcag	acatcacctc	tctgaccatc	agtttcatct	ttcctttgat	900
tggccatgtc	acgggttttc	tgtctggcact	ttttaccac	cagtcttggc	aaagggtgcag	960
gacaatttcc	ttagaaactg	gagctcagaa	tattcagatg	tgcataccca	tgtctccagtt	1020
atcttttact	gctgagcact	tgggtccagat	gttgagtctt	ccactggcct	atggactctt	1080
ccagctgata	gatggatttc	ttattgttgc	agcatatcag	acgtacaaga	ggagattgaa	1140
gaacaaacat	ggaaaaaaga	actcagggtg	cacagaagtc	tgccatacga	ggaaatcgac	1200
ttcttccaga	gagaccaatg	ccttcttggg	gggtgaatgaa	gaagggtgcca	tcactcctgg	1260
gccaccaggg	ccaatggatt	gccacagggc	tctcgagcca	gttggccaca	tcacttcatg	1320
tgaatagcag	ggactagctg	gctggactgg	cccccttctt	tttcagtggc	cagtaaagac	1380
agtgtgcagc	tgacacatga	atcttgttgg	tagggccagt	gtgaatattt	aagtgttcaa	1440
tgttagaata	tttatatttt	catgtggatt	gtgaattgtg	atgggatcac	ttttggagat	1500
tcccatttca	gggagtttct	tctgggggtt	aacataacgt	atcaatgagc	tgccttgtat	1560
gttcttgtct	tcccatgcag	caaatagtag	cctggcccta			1600

<210> 4

<211> 1317

<212> DNA

<213> Homo sapiens

<400> 4

atgattagaa	aactttttat	tgttctactt	ttgttgcttg	tgactataga	agaagcaagg	60
atgtcatcgc	tcagttttct	gaatatagag	aagactgaaa	tactattttt	cacaaagact	120
gaagaaacca	tccttgtaag	ttcaagctac	gaaaataaac	ggcctaattc	cagccacctc	180
tttgtgaaaa	tagaagatcc	taaaatacta	caaattggtga	atgtggccaa	gaagatctca	240
tcagatgcta	caaactttac	cataaatctg	gtgactgatg	aagaaggaga	aacaaatgtg	300
actattcaac	tctgggattc	tgaaggtagg	caagaaagac	tcattgaaga	aatcaagaat	360
gtgaaagtca	aagtgtctca	acaaaaagac	agtctactcc	aggcaccaat	gcataattgat	420
agaaayatcc	taatgcttat	tttaccacta	atactattga	ataagtgtgc	atttggttgt	480
aagattgaat	tacagctgtt	tcaaacagta	tgggaagagac	ctttgccagt	aattcttggg	540
gcagttacac	agttttttct	gatgccattt	tgcgggtttc	ttttgtctca	gattgtggca	600
ttgcctgagg	cgcaagcttt	tggagttgta	atgacctgca	cgtgcccagg	agggggtggg	660
ggctatctct	ttgctctgct	tctagatgga	gatttcacat	tggccatttt	gatgacttgc	720
acatcaacat	tattggctct	gatcatgatg	cctgtcaatt	cttatatata	cagtaggata	780
ttagggttgt	caggtacatt	ccatattcct	gtttctaaaa	ttgtgtcaac	actccttttc	840
atacttgtgc	cagtatcaat	tggaaatagtc	atcaagcata	gaataacctga	aaaagcaagc	900
ttcttagaga	gaataaattag	acctctgagt	tttatatttaa	tgttcgtagg	aattttatttg	960
actttcacag	tgggattagt	gttcttaaaa	acagataatc	tagagggtgat	tctgttgggt	1020
ctcttagttc	ctgctttggg	tttgctgttt	gggtactcct	ttgctaaagt	ttgtacgctg	1080
cctcttctctg	tttgtaaaac	tgttgctatt	gaaagtggga	tggttaaata	tttcttagct	1140
cttgccgtta	ttcagctgtc	ttttccacag	tccaaggcca	atttagcttc	tgtggctcct	1200
tttacagtag	ccatgtgttc	tggatgtgaa	atgttactga	tcattctagt	ttacaaggct	1260
aagaaaagat	gtatcttttt	cttacaagat	aaaaggaaaa	gaaatttctc	aatctaa	1317

<210> 5

<211> 438

<212> PRT

WO 02/072774

PCT/US02/07438

&lt;213&gt; Homo sapiens

&lt;400&gt; 5

```

Met Ile Arg Lys Leu Phe Ile Val Leu Leu Leu Leu Val Thr Ile
 1           5           10           15
Glu Glu Ala Arg Met Ser Ser Leu Ser Phe Leu Asn Ile Glu Lys Thr
          20           25           30
Glu Ile Leu Phe Phe Thr Lys Thr Glu Glu Thr Ile Leu Val Ser Ser
          35           40           45
Ser Tyr Glu Asn Lys Arg Pro Asn Ser Ser His Leu Phe Val Lys Ile
          50           55           60
Glu Asp Pro Lys Ile Leu Gln Met Val Asn Val Ala Lys Lys Ile Ser
65           70           75           80
Ser Asp Ala Thr Asn Phe Thr Ile Asn Leu Val Thr Asp Glu Glu Gly
          85           90           95
Glu Thr Asn Val Thr Ile Gln Leu Trp Asp Ser Glu Gly Arg Gln Glu
          100          105          110
Arg Leu Ile Glu Glu Ile Lys Asn Val Lys Val Lys Val Leu Lys Gln
          115          120          125
Lys Asp Ser Leu Leu Gln Ala Pro Met His Ile Asp Arg Asn Ile Leu
          130          135          140
Met Leu Ile Leu Pro Leu Ile Leu Leu Asn Lys Cys Ala Phe Gly Cys
145          150          155          160
Lys Ile Glu Leu Gln Leu Phe Gln Thr Val Trp Lys Arg Pro Leu Pro
          165          170          175
Val Ile Leu Gly Ala Val Thr Gln Phe Phe Leu Met Pro Phe Cys Gly
          180          185          190
Phe Leu Leu Ser Gln Ile Val Ala Leu Pro Glu Ala Gln Ala Phe Gly
          195          200          205
Val Val Met Thr Cys Thr Cys Pro Gly Gly Gly Gly Gly Tyr Leu Phe
          210          215          220
Ala Leu Leu Leu Asp Gly Asp Phe Thr Leu Ala Ile Leu Met Thr Cys
225          230          235          240
Thr Ser Thr Leu Leu Ala Leu Ile Met Met Pro Val Asn Ser Tyr Ile
          245          250          255
Tyr Ser Arg Ile Leu Gly Leu Ser Gly Thr Phe His Ile Pro Val Ser
          260          265          270
Lys Ile Val Ser Thr Leu Leu Phe Ile Leu Val Pro Val Ser Ile Gly
          275          280          285
Ile Val Ile Lys His Arg Ile Pro Glu Lys Ala Ser Phe Leu Glu Arg
          290          295          300
Ile Ile Arg Pro Leu Ser Phe Ile Leu Met Phe Val Gly Ile Tyr Leu
305          310          315          320
Thr Phe Thr Val Gly Leu Val Phe Leu Lys Thr Asp Asn Leu Glu Val
          325          330          335
Ile Leu Leu Gly Leu Leu Val Pro Ala Leu Gly Leu Leu Phe Gly Tyr
          340          345          350
Ser Phe Ala Lys Val Cys Thr Leu Pro Leu Pro Val Cys Lys Thr Val
          355          360          365
Ala Ile Glu Ser Gly Met Leu Asn Ser Phe Leu Ala Leu Ala Val Ile
          370          375          380
Gln Leu Ser Phe Pro Gln Ser Lys Ala Asn Leu Ala Ser Val Ala Pro
385          390          395          400
Phe Thr Val Ala Met Cys Ser Gly Cys Glu Met Leu Leu Ile Ile Leu
          405          410          415
Val Tyr Lys Ala Lys Lys Arg Cys Ile Phe Phe Leu Gln Asp Lys Arg

```



WO 02/072774

PCT/US02/07438

	420	425	430			
Lys Arg Asn Phe Leu Ile						
435						
<210> 6						
<211> 1777						
<212> DNA						
<213> Homo sapiens						
<400> 6						
gcataagcag	aaatgaggaa	atcaagagga	agagattaga	tttctgttgt	gataaatcga	60
atctgttaaa	tgccatgact	ttttaattgt	cttaatcaca	agttaaaccg	gttgtgttgc	120
tgcttagatg	gctatatatt	tgtttaaaag	tacagcagtc	cctcctactg	gactttgatc	180
ctacaaaaac	aactgttatc	taactcacc	tcagactgtc	actggaacac	ctgcatgaag	240
aatgttcttt	cattttttta	aaacgatttt	gcataatga	tttatttcag	ctttcaaaat	300
gattagaaaa	cttttttattg	ttctactttt	gttgcttgtg	actatagaag	aagcaaggat	360
gtcatcgctc	agtttttctga	atatagagaa	gactgaaata	ctattttttca	caaagactga	420
agaaaccatc	cttgtaagtt	caagctacga	aaataaacgg	cctaattcca	gccacctctt	480
tgtgaaaata	gaagatccta	aaatactaca	aatggtgaat	gtggccaaga	agatctcatc	540
agatgctaca	aactttacca	taaatctggt	gactgatgaa	gaaggagaaa	caaagtgtgac	600
tattcaactc	tgggattctg	aaggtaggca	agaaagactc	attgaagaaa	tcaagaatgt	660
gaaagtcaaa	gtgctcaaac	aaaaagacag	tctactccag	gcaccaatgc	atattgatag	720
aaayatccta	atgcttattt	taccactaat	actattgaat	aagtgtgcat	ttggttgtaa	780
gattgaatta	cagctgtttc	aaacagtatg	gaagagacct	ttgccagtaa	ttcttggggc	840
agttacacag	ttttttctga	tgccattttg	cgggtttctt	ttgtctcaga	ttgtggcatt	900
gcctgaggcg	caagcttttg	gagttgtaat	gacctgcacg	tgcccaggag	ggggtggggg	960
ctatctcttt	gctctgcttc	tagatggaga	tttcacattg	gccattttga	tgacttgcac	1020
atcaacatta	ttggctctga	tcatgatgcc	tgtcaattct	tatatataca	gtaggatatt	1080
agggttgtca	ggtacattcc	atattcctgt	ttctaaaatt	gtgtcaacac	tccttttcat	1140
acttgtgcc	gtatcaattg	gaatagtc	caagcataga	atacctgaaa	aagcaagctt	1200
cttagagaga	ataattagac	ctctgagttt	tatttttaatg	ttcgtaggaa	tttatttgac	1260
tttcacagtg	ggattagtgt	tcttaaaaac	agataatcta	gaggtgattc	tgttgggtct	1320
cttagttcct	gctttgggtt	tgctgttttg	gtactccttt	gctaaagttt	gtacgctgcc	1380
tcttcctgtt	tgtaaaactg	ttgctattga	aagtgggatg	ttaaatagtt	tcttagctct	1440
tgccgttatt	cagctgtctt	ttccacagtc	caaggccaat	ttagcttctg	tggctccttt	1500
tacagtagcc	atgtgttctg	gatgtgaaat	gttactgac	attctagtgt	acaaggctaa	1560
gaaaagatgt	atctttttct	tacaagataa	aaggaaaaga	aatttcctaa	tctaacaatt	1620
aaagcattac	tgaattccta	ctctgggtag	ggcactgtgg	gagagtaaaa	gaataactat	1680
aatgatacct	gctcttaagc	cacttataat	gtaattaaaa	atctgacatt	ggggggtaga	1740
gaatacatat	atatatatac	acacacacac	acacaca			1777